arXiv:1302.1959v1 [math-ph] 8 Feb 2013

Entanglement Entropy for a Coupled Harmonic Oscillator

Chisanupong Puttartrom*, Monsit Tanasittikosol†, Sikarin Yoo-Kong‡
Theoretical and Computational Physics (TCP) Group, Department of Physics,
Faculty of Science, King Mongkut's University of Technology Thonburi, Thailand, 10140.

We investigated the entanglement for a simple model of two-particles system in the harmonic confinement. The Feynman path integral approach was applied to compute the density matrix of the system. The spatial entanglement of ground-state was study using the linear entropy. We found the spatial entanglement increases while decreasing of the confining potential ω as this implies that the spatial separation between the particles increases.

PACS numbers:

I. INTRODUCTION

Quantum theory is one of the most important branches in physics and is well know for being counter-intuitive and its bizarre phenomena (nothing like we experience in everyday life). Quantum entanglement, which is the direct consequence of the EPR paradox [1], is a very unique phenomenon and has no classical counterparts. Since the discovery of quantum entanglement, a wide range of applications has been proposed, i.e., quantum cryptography, quantum teleportation and superdense coding [2]. These applications indicate that entanglement could be exploited to preform the impossible tasks which cannot be done by classical computers. Here is another new branch of physics, which is called quantum information and computing, has been born by joining quantum physics and theoretical computer science. In this respect many solid state systems, especially quantum dots, have been proposed as promising hardwares to perform quantum information processors [3]. The main difficulty of studying many-body solid state system is the complexity. The approximation methods must be introduced to obtain physical properties, see [4].

In this paper, we study the entanglement [4] of the ground state of two-body systems using Feynman path integral [5]. We will concentrate on the entanglement generated by the spatial degrees of freedom and use a coupled harmonic oscillator as a toy model. The Feynman path integral method will be used to compute the density matrix of the composite system (bipartite system). This provides the way to quantify the entanglement by calculating the linear entropy [6].

The organisation of this paper is the following. In Sect. II we introduce the toy model: a coupled harmonic oscillator and how to compute the density matrix. Then, the linear entropy will be briefly discussed and will be used to quantify the spatial entanglement of the system in Sect. III. And the last section will be devoted to summary and discussions.

II. THE SYSTEM



FIG. 1: The system of a coupled harmonic oscillator.

Consider the coupled harmonic oscillator as show in Fig. 1. The particle with mass m is trapped in the harmonic potential with frequency ω. The particle with mass M vibrates around its equilibrium with angular frequency Ω. The interaction between these two particles is modeled by Hooke's law and κ denotes the strength of the interaction. The Lagrangian is given by

$$\begin{aligned}\mathcal{L}_{x,y} &= \frac{m}{2}\dot{x}^2(\tau) - \frac{m}{2}\omega^2 x^2(\tau) + \frac{M}{2}\dot{y}^2(\tau) - \frac{M}{2}\Omega^2 y^2(\tau) \\ &\quad - \frac{\kappa}{2}\mid x(\tau) - y(\tau) \mid^2 , \qquad (1)\end{aligned}$$

where $x(\tau)$ is the position of the mass m, $y(\tau)$ is the position of the mass M and τ is time. The propagator of the composite system is then given by

$$\begin{aligned}\mathcal{K}_{x,y} &= \mathcal{K}(x(t), y(t), t; x(0), y(0), 0) \\ &= \int_{x(0)}^{x(t)} \mathcal{D}(x) \int_{y(0)}^{y(t)} \mathcal{D}(y) \mathrm{e}^{\frac{\mathrm{i}}{\hbar}\mathcal{S}} , \qquad (2)\end{aligned}$$

where the action is given by

$$S = \int_0^t \mathrm{d}t \mathcal{L}_{x,y} . \qquad (3)$$

We consider the case of the imaginary time: $t \to -\mathrm{i}\beta$ where $\beta = 1/k_{\mathrm{B}}T$ and T is temperature and k_{B} is Boltzmann constant [7]. The propagator Eq. (2) becomes the density matrix.

$$\rho_{x,y} = \int_{x(0)}^{x(\beta)} \mathcal{D}(x) \int_{y(0)}^{y(\beta)} \mathcal{D}(y) \mathrm{e}^{-S_\beta} , \qquad (4)$$

---

*chisanupong58@gmail.com
†monsit.tan@kmutt.ac.th
‡sikarin.yoo@kmutt.ac.th

where

$$S_\beta \;=\; \int_0^\beta \mathrm{d}\tau \mathcal{L}_{x,y} \; . \qquad (5)$$

The reduced density matrix can be obtained by tracing out the variable y of mass M [8].

$$\begin{aligned} \rho_{\rm red} \equiv \rho_x \;&=\; \mathrm{Tr}_{\rm y} \rho_{x,y} = \int_{x(0)}^{x(\beta)} \mathcal{D}(x) \int_y^y \mathcal{D}(y) e^{-S_\beta} \\ &=\; \left(\frac{m}{4\pi\hbar \triangle(0)} \right)^{\frac{1}{2}} \left(\frac{\sinh(\frac{\Omega_{\rm eff}\beta}{2})}{\sinh(\frac{z_+\beta}{2})\sinh(\frac{z_-\beta}{2})} \right) \\ &\quad \times \mathrm{e}^{[\frac{m\ddot{\triangle}(0)}{4}(x(\beta)-x(0))^2 - \frac{m}{4\triangle(0)}(x(\beta)+x(0))^2]} \; , \end{aligned} \qquad (6)$$

where

$$\begin{aligned} \triangle\,(\tau) \;&=\; \frac{{z_+}^2 - {\Omega^2}_{\rm eff}}{{z_+}^2 - {z_-}^2} \frac{\cosh(z_+(\tau - \frac{\beta}{2}))}{z_+ \sinh(z_+ \sinh(\frac{z_+\beta}{2}))} \\ &\quad + \frac{{z_-}^2 - {\Omega^2}_{\rm eff}}{{z_-}^2 - {z_+}^2} \frac{\cosh(z_-(\tau - \frac{\beta}{2}))}{z_- \sinh(z_- \sinh(\frac{z_-\beta}{2}))} \; , \end{aligned} \qquad (7a)$$

$$2z_\pm^2 \;=\; (\Omega_{\rm eff}^2 + \frac{\kappa}{m}) \pm \sqrt{(\Omega_{\rm eff}^2 + \frac{\kappa}{m})^2 - 4\omega^2\Omega_{\rm eff}^2} \; , \qquad (7b)$$

$${\Omega^2}_{\rm eff} \;=\; \Omega^2 + \frac{\kappa}{M} \; . \qquad (7c)$$

III. LINEAR ENTROPY

The entanglement generated by the continuous spatial degree of freedom is expected to depend on the interplay between the strength of the confining potential and interaction. Then we are interested to investigate how the spatial entanglement changes with the system parameter.

The linear entropy of the reduced density matrix

$$\mathcal{S}_{\rm L} \;=\; 1 - \mathrm{Tr}\rho_{\rm red}^2 \; , \qquad (8)$$

where $0 \le \mathcal{S}_{\rm L} \le 1$ is used to measure the entanglement for a pure state and continuous case. For the continuous variable case, $\rho_{\rm red}^2$ can be computed by

$$\rho_{\rm red}^2 = \int \mathrm{d}x \rho_{\rm red}(x,x')\rho_{\rm red}(x',x) \; , \qquad (9)$$

and

$$\mathrm{Tr}\rho_{\rm red}^2 = \int \mathrm{d}x \rho_{\rm red}^2(x,x) \; . \qquad (10)$$

The system would be completely disentangled where $\mathcal{S}_{\rm L} = 0$. On the other hand, the system would be maximally entangled state if $\mathcal{S}_{\rm L} = 1$.

The linear entropy of our systems is

$$\mathcal{S}_{\rm L} = 1 - \left(\frac{\mathrm{e}^{\Omega_{\rm eff}\beta}}{\mathrm{e}^{\beta(\mathrm{z}_+ + \mathrm{z}_-)}} \right) \left(\frac{A}{B} \right)^{\frac{1}{2}} \; , \qquad (11)$$

where

$$A \;=\; \left[\left(\Omega_{\rm eff}^2 + \frac{\kappa}{m} \right)^2 - 4\omega^2\Omega_{\rm eff}^2 \right] \omega\Omega_{\rm eff} \; , \qquad (12a)$$

$$\begin{aligned} B \;&=\; \omega\Omega_{\rm eff} \left(\frac{\kappa}{m} - \Omega_{\rm eff}^2 \right)^2 \\ &\quad - \left(\Omega_{\rm eff}^2 + \frac{\kappa}{m} + 2\omega\Omega_{\rm eff} \right) \left(\omega^2\Omega_{\rm eff}^2 - \frac{\kappa}{m}\Omega_{\rm eff}^2 \right) \end{aligned} \qquad (12b)$$

To obtain Eq. (11), we take the limit where $\beta \to \infty$ in order to ensure that the system is the ground state. In this limit, the system is maximally entangled state when $\mathcal{S}_{\rm L} = 1$ leading to the condition that is

$$\omega^2 = \frac{(\Omega^2 + \kappa/\mu)^2}{4(\Omega^2 + \kappa/M)} \equiv \omega_{\rm M}^2 \; , \qquad (13)$$

where

$$\frac{1}{\mu} = \frac{1}{M} + \frac{1}{m} \; .$$

In the case of the separable state, $\mathcal{S}_{\rm L} = 0$, we obtain the condition

$$\omega^2 = \frac{(\Omega^2 + \kappa/M)}{16} \equiv \omega_{\rm S}^2 \; . \qquad (14)$$

Eq. (13) and Eq. (14) show that the confining potential strength ω (external parameter) depends on the parameter Ω and κ for the case of maximally entangled and the case of separable state, respectively.

Let us, we consider for the case that Ω becomes very large. We obtain

$$\omega_{\rm M}^2 \simeq \tfrac{\Omega^2}{4} \text{ and } \omega_{\rm S}^2 \simeq \tfrac{\Omega^2}{16}$$

Obviously, the confining parameter ω solely depends on the strength of the variable Ω. On the other hand, if Ω becomes infinitesimally small, we obtain

$$\omega_{\rm M}^2 \simeq \tfrac{\kappa M}{4\mu^2} \text{ and } \omega_{\rm S}^2 \simeq \tfrac{\kappa}{16M}$$

In this case we see that confining parameter depends on the strength of the interaction. From these results we can adjust the entanglement between particle mass m and particle mass M by tuning the strength of ω with vibrating frequency Ω and coupling constant κ.



FIG. 2: Linear entropy versus ω



FIG. 3: Linear entropy versus Ω_{eff}

In Fig. 2 shows the numerical results of how $\mathbb{S}_{\text{L}}$ changes with respect to ω. It is clear the spatial entanglement increases when the confining potential ω decreases. This is because the spatial separation between m and M increases due to low confining potential and hence $\mathbb{S}_{\text{L}}$ increases [4]. In Fig. 3, the confining potential ω is fixed while the interaction strength κ is varied. It is found that the spatial entanglement increases with the strength of the interaction.

IV. CONCLUSION

We have studied the spatial entanglement through the linear entropy for the coupled harmonic oscillator. The density matrix of the system could be exactly computed by using Feynman path integral approach. We obtain the condition on system parameters for maximally entangled and completely disentangled state. The system in Sect. II can be extended by adding more of particles with mass M. Then this can be viewed as the particles with M constitute the lattice structure while the particle with mass m is the charge moving through the lattice. The spatial entanglement between the charge (particle with mass m) and the lattice (particle with mass M) could possibly be studied using the same technique in this paper. Furthermore, this method could also possible be used to study the entanglement in polaron systems.



FIG. 4: The moving particle in the lattice.

---

[1] A. Einstein, A; B. Podolsky, N. Rosen, Can Quantum-Mechanical Description of Physical Reality be Considered Complete?, Phys Rev. 47 (10), pp.777-780, (1935) ; J.S. Bell, On the Einstein Poldolsky Rosen paradox, Physics 1,195-200, (1964).

[2] C. H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, W. K. Wootters, Teleporting an Unknown Quantum State via Dual Classical and Einstein-Podolsky-Rosen Channels, Phys. Rev. Lett. 70, pp.1895-1899, (1993); M. A. Nielsen, I. L. Chuang, Quantum Computation and Quantum Information, Cambridge University Press (2000); Hoi-Kwong Lo, S. Popescu, T. Spiller, Introduction To Quantum Computation and Information, World Scientific Publishing Co. Pte. Ltd; D. Bouwmeester, A. Ekert, A. Zeilinger, The Physics of Quantum Information, Springer.

[3] E. Biolatti, et al, Electro-optical properties of semiconductor quantum dots: Application to quantum information processing, , Phys Rev. B 65, 075300, (2002); D. Loss, D. P. DiVincenzo, Quantum computation with quantum dots, Phys Rev. A 57, 120, (1998); T. Tanamoto, Quantum gates by coupled asymmetric quantum dots and controlled-NOT-gate operation Phys Rev. A 61, 022305, (2000).

[4] K. Capella, A bird's-eye view of density-functional theory, Brazilian Journal of physic 36, 1318,2006; J. P. Coe, A. Sudbery and I. D'Amico, Entanglement in GaAs and CdSe quantum dots: Exact calculation and DFT approximations, Microelectronics Journal 40 (2009), pp.449–501.

[5] Richard P. Feynman, Albert R. Hibbs, Quantum Mechanics and Path Integrals: Emended Edition , McGraw-Hill, (1965); C. Grosche, F. Steiner, Handbook of Feynman Path Integrals, Springer, (1998);

[6] Thomas Brumme, Entanglement in Continuous Degrees of Freedom, Max Planck Institute for the Physics of Complex System, Dresden, (2006); W.J. Munro, D. F. V. James, A. G. White and P. G. Kwiat, Maximizing the entanglement two mixed qubits, Phys Rev A 64, 030302, (2001)

[7] Richard P. Feynman, Statistical Mechanics A set of Lectures, W.A. Benjamin, Inc., (1972)

[8] Adamocski, J., Gerach, B., Leschke, Explocit Evaluation of Certain Gaussian Functional Integrals Arising in Problems of Statistical Physics, J. Math. Phy. 23, pp.243-249, (1982); Cheng, B.K., Path Integration of the Harmonic Oscillator with a Two-Time Quadratic Action, Phy.Lett. A 103, pp.-357-362, (1984); Sikarin Yoo-Kong, The propagator for a charged particle in an electromagnetic field and a series of non-local harmonic oscillators, Journal-Korean Physical Society 49(5), pp.1998-2000, (1998)